Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Journal of Organometallic Chemistry 669 (2003) 211
Journal of Organometallic Chemistry
www.elsevier.com/locate/jorganchem

Author Index of Volume 669

Abd-Elzaher, M.M., 6
Ali, S., 72
Angeles Paz-Sandoval, M., 210
Avent, A.G., 87

Bassindale, A.R., 154
Baukov, Y.I., 154
Beckmann, J., 149
Berke, H., 14
Bhatti, M.H., 72
Borbaruah, M., 154

Cariati, E., 44
Caubet, A., 164
Cebulski, A., 64
Chai, J.-F., 57
Chang, F.-C., 182
Chen, J.-D., 182
Coles, M.P., 209

Dakternieks, D., 149
Dötz, K.H., 1
Dragonetti, C., 44
Duthie, A., 149

Eng, G., 109
Esther Sánchez-Castro, M., 210

Fischer, H., 6
Font-Bardía, M., 164
Fox, T., 14

Ghebreyessus, K.Y., 48
Glynn, S.J., 154
Gordillo-Román, B., 210
Graziani, O., 200

Haase, T., 25
Hamon, J.-R., 200
Harfoot, G.J., 106
Heeg, M.J., 37
Hernández-Martínez, J., 210
Hidai, M., 124
Hillier, A.C., 14
Hitchcock, P.B., 87, 101

Janes, E., 1
Jerzykiewicz, L., 64
Juárez-Saavedra, P., 210

Kim, Y., 32
Kohse-Höinghaus, K., 25
Krishnamurthy, S.S., 79
Kubo, K., 141
Kumar, A., 109
Kunicki, A.R., 64

Lang, H., 25
Leigh, G.J., 87, 101, 207, 208, 209
Leschke, M., 25
Liang, H.-C., 182
López, C., 164
Lucenti, E., 44

Madura, I., 64
Maisel, H.E., 72
Mathieu, R., 172
Matsubayashi, G.-e., 141
Milius, W., 72
Mizobe, Y., 124
Monari, M., 135

Nagao, S., 124
Nakamoto, M., 141
Nakano, M., 141
Nath, M., 109
Negrebetsky, V.V., 154
Nelson, J.H., 48
Nethaji, M., 79
Nicholson, B.K., 106

Ogino, H., 189
Okazaki, M., 189

Palazzi, A., 135
Parker, D.J., 154
Pokharia, S., 109

Raghuraman, K., 79
Roberto, D., 44
Román Torres-Lubián, J., 210

Sato, T., 189
Schmalle, H.W., 14
Schmidt, H., 25
Seino, H., 124
Selva, S., 135
Shen, Y., 25
Smith, N.A., 149
Solans, X., 164
Solsona, A., 172
Song, X., 109
Stagni, S., 135
Suades, J., 172
Szumacher, S., 64

Tamura, H., 141
Tang, L.-F., 57
Taylor, P.G., 154
Tilset, M., 200
Tobita, H., 189
Togrou, M., 87, 101
Toupet, L., 200
Turtle, R., 154

Verkade, J.G., 32

Wang, J.-C., 182
Wang, J.-T., 57
Weibert, B., 6
Winter, C.H., 37
Wrackmeyer, B., 72
Wu, Y.-Y., 182

Xia, A., 37

Yang, P.-Y., 182

Zachara, J., 64
Zhao, S.-B., 57

Elsevier Science B.V.
doi:10.1016/S0022-328X(03)00084-6

Available online at www.sciencedirect.com
SCIENCE DIRECT®
ELSEVIER
Journal of Organometallic Chemistry 669 (2003) 213–216
Journal of Organometallic Chemistry
www.elsevier.com/locate/jorganchem

Subject Index of Volume 669

Acetonitrile
Borohydride reduction of a rhenium-bound acetonitrile: an example of a chelating iminoborane ligand at a low valent metal center, 14
Acyl complex
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Addition
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes, 6
Aldol condensation
Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand, 1
Alkenes
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72
Alkoxy ligands
Influence of alkoxy ligands on the Cp–Al bonding mode in $[Cp_2Al\text{-}\mu\text{-}OR]_2$ from X-ray crystallographic and ^{27}Al-NMR spectroscopic solution studies, 64
Alkyl complex
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Alkyne complexes
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes, 6
Aminate complexes
Synthesis of titanium(IV) (cyclopentadienylalkyl)aminate complexes, 101
Amino acid
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109
Amphiphilic phosphines
Hydridorhodium(I) complexes with amphiphilic polyether phosphines: NMR study and biphasic hydroformylation of 1-octene, 172

Bicyclic ligand
Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane, 32
Bimetallic selenido clusters
Syntheses of a series of trinuclear MIr_2 or pentanuclear MIr_4 bimetallic bis(selenido) and selenido–sulfido clusters (M = Pd, Pt, Fe, Co) from diiridium μ-bis(hydroselenido) and μ-hydroselenido–hydrosulfido complexes $[\{(\eta^5\text{-}C_5Me_5)IrCl\}_2(\mu\text{-}SeH)(\mu\text{-}EH)]$ (E = Se, S), 124
Bioorganometallic chemistry
Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand, 1
Biphasic hydroformylation
Hydridorhodium(I) complexes with amphiphilic polyether phosphines: NMR study and biphasic hydroformylation of 1-octene, 172

Borohydride
Borohydride reduction of a rhenium-bound acetonitrile: an example of a chelating iminoborane ligand at a low valent metal center, 14

C-Deprotonated-2-phenylpyridine complexes
Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of $[Au(\eta^2\text{-}C,N\text{-}ppy)(\eta^2\text{-}S,S\text{-}C_8H_4S_8)]\cdot 0.5DMF$, 141
C-glycosides
Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand, 1
Chelate
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand $\kappa^2(Si,N)$-$SiMe_2O(2\text{-}C_5H_4N)$, 189
Chiral phosphorus ligands
Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of $[Ru_3(\mu\text{-}CO)(CO)_7(\mu_3\text{-}S)\{Ph_2PN((S)\text{-}*CHMePh)PPh_2\text{-}\kappa^2P,P\}]$ and $[Rh(CO)Cl\{Ph_2PN((S)\text{-}*CHMePh)P(S)Ph_2\}\text{-}\kappa^2P,S]$: Part 16. Organometallic chemistry of diphosphazanes, 79
Chromium
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes, 6
Chromium carbene complexes
Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand, 1
Cluster
Anomalous reaction of an aryl silane with $Co_2(CO)_8$; characterisation of $Me_2NC_6H_4Si[Co(CO)_4][OCCo_3(CO)_9]_2$, 106
Cluster compounds
Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of $[Ru_3(\mu\text{-}CO)(CO)_7(\mu_3\text{-}S)\{Ph_2PN((S)\text{-}*CHMePh)PPh_2\text{-}\kappa^2P,P\}]$ and $[Rh(CO)Cl\{Ph_2PN((S)\text{-}*CHMePh)P(S)Ph_2\}\text{-}\kappa^2P,S]$: Part 16. Organometallic chemistry of diphosphazanes, 79
Cobalt carbonyl
Anomalous reaction of an aryl silane with $Co_2(CO)_8$; characterisation of $Me_2NC_6H_4Si[Co(CO)_4][OCCo_3(CO)_9]_2$, 106
Conformation
Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands, 182
Coupling constant
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154
Cp–Al bond mode
Influence of alkoxy ligands on the Cp–Al bonding mode in $[Cp_2Al\text{-}\mu\text{-}OR]_2$ from X-ray crystallographic and ^{27}Al-NMR spectroscopic solution studies, 64

Elsevier Science B.V.
doi:10.1016/S0022-328X(03)00083-4

Crystal structure
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Cubic
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium, 37
Cyclopentadienyl
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium, 37
Synthesis of titanium(IV) (cyclopentadienylalkyl)aminate complexes, 101
Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2S$} and {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2CH_2$}, 57
Cycloplatination
Synthesis, characterisation and study of the first luminescent platinum(II) compound with a $[C,N,S]^-$ terdentate ligand. X-ray crystal structure of [Pt{C_6H_4–CH=N–(C_6H_4–2-SMe)}Cl], 164

[4+2]Diels–Alder cycloaddition
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones, 48
Difunctional phosphorus ligand
Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane, 32
Diiridium hydroselenido complexes
Syntheses of a series of trinuclear MIr_2 or pentanuclear MIr_4 bimetallic bis(selenido) and selenido–sulfido clusters (M = Pd, Pt, Fe, Co) from diiridium μ-bis(hydroselenido) and μ-hydroselenido–hydrosulfido complexes [{(η^5-C_5Me_5)IrCl}$_2$(μ-SeH)(μ-EH)] (E = Se, S), 124
Dipyridylformamidine
Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands, 182
Dithiolate complexes
Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of [Au(η^2-*C*,*N*-ppy)(η^2-*S*,*S*-$C_8H_4S_8$)]·0.5DMF, 141

Electrical conductivities
Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of [Au(η^2-*C*,*N*-ppy)(η^2-*S*,*S*-$C_8H_4S_8$)]·0.5DMF, 141

Fe(II) organometallic complexes
Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative., 135
Fluorine NMR
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154

Gold(III) complexes
Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of [Au(η^2-*C*,*N*-ppy)(η^2-*S*,*S*-$C_8H_4S_8$)]·0.5DMF, 141

3*H*-Pyrazole complex
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes, 6
Hydridorhodium(I) complexes
Hydridorhodium(I) complexes with amphiphilic polyether phosphines: NMR study and biphasic hydroformylation of 1-octene, 172
Hydroboration
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72

Interannular conjugation
Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative., 135
Iron complex
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Iron complexes
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand κ^2(*Si*,*N*)-$SiMe_2O(2\text{-}C_5H_4N)$, 189
Isomerisation
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72

Lithiation reaction
The lithiation of arylamines and the preparation of cyclopentadienyltitanium(IV) arylaminate complexes, 87
Lithio derivatives
The lithiation of arylamines and the preparation of cyclopentadienyltitanium(IV) arylaminate complexes, 87
Luminiscence
Synthesis, characterisation and study of the first luminescent platinum(II) compound with a $[C,N,S]^-$ terdentate ligand. X-ray crystal structure of [Pt{C_6H_4–CH=N–(C_6H_4–2-SMe)}Cl], 164

Magnesium
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium, 37
Magnesocene
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium, 37
Metal carbonyl complexes
Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands, 182
Metallamacrocycle
Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2S$} and {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2CH_2$}, 57
Molybdenum
Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2S$} and {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2CH_2$}, 57
Multi-nuclear NMR
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109

NMR
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72
Nucleophilic substitution
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of

pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154

Organometallic compounds
Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of [Ru_3(μ-CO)(CO)$_7$(μ$_3$-S){$Ph_2PN((S)$-*CHMePh)PPh_2-κ$^2P,P$}] and [Rh(CO)Cl{$Ph_2PN((S)$-*CHMePh)P(S)Ph_2}-κ$^2P,S$]: Part 16. Organometallic chemistry of diphosphazanes, 79
Organometallic monosaccharides
Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a *C*-monosaccharide ligand, 1
Organosilanes
Anomalous reaction of an aryl silane with $Co_2(CO)_8$; characterisation of $Me_2NC_6H_4Si[Co(CO)_4][OCCo_3(CO)_9]_2$, 106
Organotin(IV)
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109

Palladium(II) chloride
Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane, 32
Pentacoordinate
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154
Peptide
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109
Phosphole
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones, 48
Photochemical decarbonylation
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Photolysis
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand κ^2(*Si*,*N*)-$SiMe_2O$(2-C_5H_4N), 189
Platinum(II)
Synthesis, characterisation and study of the first luminescent platinum(II) compound with a [C,N,S]$^-$ terdentate ligand. X-ray crystal structure of [Pt{C_6H_4–CH=N–(C_6H_4–2-SMe)}Cl], 164
P,S-donor ligands
Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of [Ru_3(μ-CO)(CO)$_7$(μ$_3$-S){$Ph_2PN((S)$-*CHMePh)PPh_2-κ$^2P,P$}] and [Rh(CO)Cl{$Ph_2PN((S)$-*CHMePh)P(S)Ph_2}-κ$^2P,S$]: Part 16. Organometallic chemistry of diphosphazanes, 79

Rhenium
Borohydride reduction of a rhenium-bound acetonitrile: an example of a chelating iminoborane ligand at a low valent metal center, 14
Ruthenium
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones, 48
Ruthenium carbonyl clusters synthesis
Reproducible high-yield syntheses of [$Ru_3(CO)_{12}$], [$H_4Ru_4(CO)_{12}$], and [$Ru_6C(CO)_{16}]^{2-}$ by a convenient two-step methodology involving controlled reduction in ethylene glycol of $RuCl_3 \cdot n$-H_2O, 44
Ruthenium trichloride reductive carbonylation
Reproducible high-yield syntheses of [$Ru_3(CO)_{12}$], [$H_4Ru_4(CO)_{12}$], and [$Ru_6C(CO)_{16}]^{2-}$ by a convenient two-step methodology involving controlled reduction in ethylene glycol of $RuCl_3 \cdot n$-H_2O, 44

Secondary bonding
Secondary bonding in *para*-substituted diphenyltellurium dichlorides (p-$XC_6H_4)_2TeCl_2$ (X = H, Me, MeO) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of (p-$MeC_6H_4)_2TeCl_2$, 149
Silanes
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72
Silicon
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154
Silyl complexes
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand κ^2(*Si*,*N*)-$SiMe_2O$(2-C_5H_4N), 189
^{119}Sn-Mössbauer
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109
Structural correlation
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154
Structure–activity relationship
Comparative study of structure–activity relationship of di- and triorganotin(IV) derivatives of amino acid and peptides, 109

Tellurium
Secondary bonding in *para*-substituted diphenyltellurium dichlorides (p-$XC_6H_4)_2TeCl_2$ (X = H, Me, MeO) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of (p-$MeC_6H_4)_2TeCl_2$, 149
^{125}Te MAS NMR spectroscopy
Secondary bonding in *para*-substituted diphenyltellurium dichlorides (p-$XC_6H_4)_2TeCl_2$ (X = H, Me, MeO) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of (p-$MeC_6H_4)_2TeCl_2$, 149
Tetrazolate ligands
Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative., 135
Thiolato
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium, 37
Tin
Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of {p-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2S$} and {$p$-[$(CO)_3MoC_5H_4$-$C(O)]_2C_6H_4$}{$(Ph_2Sn)_2CH_2$}, 57
Titanium
Synthesis of titanium(IV) (cyclopentadienylalkyl)aminate complexes, 101
Transfer hydrogenation
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones, 48
Trinuclear and pentanuclear cores
Syntheses of a series of trinuclear MIr_2 or pentanuclear MIr_4 bimetallic bis(selenido) and selenido–sulfido clusters (M = Pd, Pt, Fe, Co) from diiridium μ-bis(hydroselenido) and μ-hydro-

selenido–hydrosulfido complexes [{(η^5-C_5Me_5)IrCl}$_2$(μ-SeH)(μ-EH)] (E = Se, S), 124
Tris(pyrazolyl)borate
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex, 200
Tungsten
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3*H*-pyrazole complexes, 6
Tetranuclear heterodimetallic metallamacrocyles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2S$} and {*p*-[$(CO)_3MoC_5H_4C(O)]_2C_6H_4$}{$(Ph_2Sn)_2CH_2$}, 57

Vinylphosphine
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones, 48

X-ray
Hydroboration of alkyn-1-yl(methyl)silanes bearing functional substituents at silicon, 72
X-ray crystallography
Anomalous reaction of an aryl silane with $Co_2(CO)_8$; characterisation of $Me_2NC_6H_4Si[Co(CO)_4][OCCo_3(CO)_9]_2$, 106
Influence of alkoxy ligands on the Cp–Al bonding mode in [Cp_2Al-μ-OR]$_2$ from X-ray crystallographic and ^{27}Al-NMR spectroscopic solution studies, 64
Secondary bonding in *para*-substituted diphenyltellurium dichlorides (p-$XC_6H_4)_2TeCl_2$ (X = H, Me, MeO) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of (p-$MeC_6H_4)_2TeCl_2$, 149
X-ray crystal structures
Preparation and oxidation of polarized Au(III) complexes having both the *C*-deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of [Au(η^2-*C*,*N*-ppy)(η^2-*S*,*S*-$C_8H_4S_8$)]·0.5DMF, 141
X-ray diffractometry
Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative., 135
X-ray structure
Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane, 32
Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands, 182
X-ray structure analysis
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand κ^2(*Si*,*N*)-$SiMe_2O(2\text{-}C_5H_4N)$, 189
X-ray structures
The lithiation of arylamines and the preparation of cyclopentadienyltitanium(IV) arylaminate complexes, 87
X-ray study
Modelling nucleophilic substitution at silicon using solution ^{19}F-NMR chemical shift, $^1J_{Si-F}$ and $^2J_{C-F}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures, 154